



# **ALFAMAX J25**

Class: Concentrated Dairy Feeds

Registration: V20381 Wet/Act 36 of 1947

| COMPOSITION     |                         |      |      |
|-----------------|-------------------------|------|------|
| Protein         | 120                     | g/kg | Min. |
| NPN             | (Max. 28.7% of protein) |      |      |
| Moisture        | 120                     | g/kg | Max. |
| Fat             | 25                      | g/kg | Min. |
| Fat             | 100                     | g/kg | Max. |
| Fibre           | 120                     | g/kg | Max. |
| Calsium (Ca)    | 8                       | g/kg | Min. |
| Calsium (Ca)    | 15                      | g/kg | Max. |
| Phosphorus (P)  | 5                       | g/kg | Min. |
| Urea            | 12                      | g/kg | Max. |
| NETT MASS ED ka |                         |      |      |

### INGREDIENT STATEMENT

May Contain: Grain Products, Plant Protein Products, Animal Protein Products, Processed Grain By-Products, Oils & Fats, Vitamins & Minerals.

#### FEEDING RECOMMENDATIONS

Feed 400 - 500 g per litre of milk produced. Feed with roughage and/or pastures. The quantity of concentrate will depend on:

- 1) the quality of the roughage;
- 2) milk production;
- 3) stage of lactation.

## WARNING:

THIS FARM FEED MAY CONTAIN A NPN SOURCE AND / OR REGISTERED STOCK REMEDIES. FEED STRICTLY ACCORDING TO INSTRUCTIONS. SEE ATTACHED STOCK REMEDY LABELS FOR FURTHER DETAIL.

Registration Holder QUANTUM FOODS (PTY) LTD

Reg. No. 2012/124966/07 PO Box 1183, WELLINGTON, 7654 Tel.: (021) 864 8600

Batch no.:

Expiry date:

Date of manufacture:

Contains genetically modified organisms

### NPN WARNING

- VINEGAR IS AN EFFECTIVE REMEDY AGAINST NPN POISONING. MIX WITH AN EQUAL AMOUNT OF WATER.
  - DOSE HALF A BOTTLE PER CALF OR LARGE SHEEP OR 2-4
- BOTTLES PER HEAD OF CATTLE. (1 BOTTLE = 750ML)

  2. PROTECT THIS FARM FEED AGAINST RAIN. NPN IS SOLUBLE
- AND ANIMALS DRINKING SUCH A SOLUTION COULD BE POISONED.
- DO NOT FEED THIS FARM FEED INDISCRIMINATELY WITH OTHER NPN CONTAINING FARM FEEDS. CONSULT AN ANIMAL SCIENTIST.
- 4. ADAPTATION: WHEN CHANGING FROM A NPN-FREE DAIRY MEAL TO A NPN-CONTAINING MEAL IT IS ADVISABLE TO FFED A 50/50 MIXTURE OVER A PERIOD OF 4-6 DAYS.
- 5. DAIRY MEAL MUST BE FED TWO OR THREE TIMES DAILY.