



HPK 50

Class: Concentrated Dairy Feeds

Registration: V23132

Wet/Act 36 of 1947

COMPOSITION			
Protein	500	g/kg	Min.
Prot. % ex NPN	31.6 % of Protein		Max
Moisture	120	g/kg	Max.
Fat	20 / 100	g/kg	Min.
Fibre	120	g/kg	Max.
Calcium (Ca)	45 / 80	g/kg	Min.
Phosphorus (P)	20	g/kg	Min.
Urea	55	g/kg	Max.
NETT MASS 50 kg			

INGREDIENT STATEMENT

Contains: Plant Protein Products, Animal Protein Products, Processed Grain By-Products, Oils & Fats, Vitamins & Minerals.

FEEDING RECOMMENDATIONS

(1) Adaptation: When changing from a NPN-free dairy meal to a NPN-containing meal it is advisable to feed a 50/50 mixture over a period of 4 - 6 days.

(2) Dairy meal must be fed two or more times daily.

(3) Mix this concentrate with maize (see mixing instructions) to obtain a dairy meal. Feed the dairy meal at approximately 350 – 500 g per litre of milk produced. The quantity of dairy meal fed will depend on: The quality and quantity of the available grazing and/or roughage; the stage of lactation and the milk and butterfat production.

Registration Holder

QUANTUM FOODS (PTY) LTD

Reg. No. 2012/124966/07

PO Box 1183, WELLINGTON, 7654

Tel.: (021) 864 8600

Batch no.:

Expiry date:

Date of manufacture:

Contains genetically modified organisms

MIXING INSTRUCTIONS

HPK 50	150 kg
Maize Meal (Prot: 75 g/kg)	600 kg
TOTAL	750 kg

COMPOSITION OF MIXTURE

Protein	150	g/kg	Min.
Prot. % ex NPN	(Max. 19.1 % of protein)		Max.
Moisture	120	g/kg	Max.
Fat	25 / 100	g/kg	Min.
Fibre	120	g/kg	Max.
Calcium (Ca)	8 / 15	g/kg	Min.
Phosphorus (P)	5	g/kg	Min.
Urea	10	g/kg	Max.

NETT MASS 50 kg

NPN WARNING

1. VINEGAR IS AN EFFECTIVE REMEDY AGAINST NPN POISONING. MIX WITH AN EQUAL AMOUNT OF WATER. DOSE HALF A BOTTLE PER CALF OR LARGE SHEEP OR 2-4 BOTTLES PER HEAD OF CATTLE. (1 BOTTLE = 750ML)
2. PROTECT THIS FARM FEED AGAINST RAIN. NPN IS SOLUBLE AND ANIMALS DRINKING SUCH A SOLUTION COULD BE POISONED.
3. DO NOT FEED THIS FARM FEED INDISCRIMINATELY WITH OTHER NPN CONTAINING FARM FEEDS. CONSULT AN ANIMAL SCIENTIST.
4. ADAPTATION: WHEN CHANGING FROM A NPN-FREE DAIRY MEAL TO A NPN-CONTAINING MEAL IT IS ADVISABLE TO FEED A 50/50 MIXTURE OVER A PERIOD OF 4-6 DAYS.
5. DAIRY MEAL MUST BE FED TWO OR THREE TIMES DAILY.