



HPC DRY COW

Class: Concentrated Dry Cow Feeds

Registration: V23510

Wet/Act 36 of 1947

COMPOSITION			
Protein	500	g/kg	Min.
Non-protein nitrogen	34.7 % of protein		Max
Moisture	120	g/kg	Max.
Fat	25	g/kg	Min.
Fat	100	g/kg	Max.
Fibre	120	g/kg	Max.
Calcium (Ca)	8	g/kg	Min.
Calcium (Ca)	15	g/kg	Max.
Phosphorus (P)	5	g/kg	Min.
Urea	10	g/kg	Max.

NETT MASS 50 kg

INGREDIENT STATEMENT

Contains: Grain products, Plant protein products, Animal protein products, Processed grain by-products, Oils and fats, Vitamins and Minerals.

FEEDING RECOMMENDATIONS

- (1) Adaptation: When changing from a NPN-free dairy meal to a NPN-containing meal it is advisable to feed a 50/50 mixture over a period of 4-6 days.
- (2) Dairy meal must be fed two or more times daily.
- (3) Mix this concentrate with maize (see mixing instructions) to obtain a dry cow meal. Feed the dry cow meal at 6.5 – 7 kg per day to far off dry cows for a period of approximately 60 days before expected calving date. The quantity of dry cow meal fed will depend on the quality and quantity of the available grazing and/or roughage and the age of the animal.

Registration Holder

QUANTUM FOODS (PTY) LTD
Reg. No. 2012/124966/07
PO Box 1183, WELLINGTON, 7654
Tel.: (021) 864 8600

Batch no.:

Expiry date:

Date of manufacture:

Best before:

Contains genetically modified organisms

WARNING:

THIS FARM FEED CONTAINS A NPN SOURCE AND / OR REGISTERED STOCK REMEDIES. FEED STRICTLY ACCORDING TO INSTRUCTIONS. SEE ATTACHED STOCK REMEDY LABELS FOR FURTHER DETAIL.

NPN WARNING

- 1. VINEGAR IS AN EFFECTIVE REMEDY AGAINST NPN POISONING. MIX WITH AN EQUAL AMOUNT OF WATER. DOSE HALF A BOTTLE PER CALF OR LARGE SHEEP OR 2-4 BOTTLES PER HEAD OF CATTLE. (1 BOTTLE = 750ML)**
- 2. PROTECT THIS FARM FEED AGAINST RAIN. NPN IS SOLUBLE AND ANIMALS DRINKING SUCH A SOLUTION COULD BE POISONED.**
- 3. DO NOT FEED THIS FARM FEED INDISCRIMINATELY WITH OTHER NPN CONTAINING FARM FEEDS. CONSULT AN ANIMAL SCIENTIST.**
- 4. ADAPTATION: WHEN CHANGING FROM A NPN-FREE DAIRY MEAL TO A NPN-CONTAINING MEAL IT IS ADVISABLE TO FEED A 50/50 MIXTURE OVER A PERIOD OF 4-6 DAYS.**
- 5. DAIRY MEAL MUST BE FED TWO OR THREE TIMES DAILY.**